

Koodana,
Phaandana,
Khelna hai
~ toh ~
**RUKNA
MANA HAI**



ALL ABOUT SELF-INFUSION

योग
Samparaq™
Connect to Empower

INFUSION INSTRUCTIONS FOR FACTORS^{1,2}

Following information is based on product package insert guidelines for infusion



1. Wash your hands with soap and water thoroughly



2. Take the contents of the pack of fridge; leave standing to bring to room temperature



3. Get the infusion kit/tray ready

(a) 2 surgical spirit/disinfection swabs for disinfection of the vials

(b) Sticky tape to keep the butterfly needle still, plaster and cotton wool

(c) Alcohol swabs for skin disinfection

(d) Syringe and tourniquet

(e) Bottle of factor (powder) and diluents (liquid)



4. Apply local anesthesia to the site of the vein





5. Getting the factor ready for infusion (reconstitution)

(a) Remove the protective covering from the short end of double-ended needle and insert the exposed needle through stopper to the diluents vial

(b) Remove the protective covering from the long end of the double-ended needle. Invert solvents (diluents) over the upright factor (powder) vials, then rapidly insert free end of the needle through the factor (powder) vial stopper at its the center. The vaccum in the vial will draw in the solvent

(c) Disconnect the two vials by removing needle from the solvent vials stopper, then remove the needle from the factor (powder) vial

(d) Roll the vial gently between the hands without shaking it, until the solution is clear. Vigorous shaking to be avoided

(e) With the filter on the syringe; draw in the reconstituted factor into the syringe

(f) Expel air out of the the filter needle (if any)

(g) Discard the needle into the sharp box and connect the syringe to the tubing of the butterfly needle



6. Put on the tourniquet, disinfect the injection site with the alcohol swab and insert the butterfly needle with the bevel facing upwards and secure it with adhesive tape. Check that the needle is in the vein by drawing little blood into the butterfly needle, and undo the tourniquet and inject fluid, checking the back flow from time to time





7. Inject slowly; the injection should last for 1-10 minutes



8. Put on adhesive bandage and clear up everything



9. Note down the date of the injection



**SELF-INFUSION GUIDANCE THROUGH
A DEMO VIDEO AND A PATIENT EDUCATION
LEAFLET**



18001034260



purna.samparq@indegene.com





Purna Samparq -Patient Support Program is run,
managed and implemented by Indegene Limited and sponsored by
Takeda Biopharmaceuticals India Pvt. Ltd. (formerly known as Baxalta Bioscience India Pvt. Ltd)



Purna Samparq is a non promotional program and no patient
identifiable data is shared with Takeda at any stage.



18001034260



purna.samparq@indegene.com

REFERENCES: 1. <https://www.hog.org/handbook/article/3/35/steps-to-prepare-and-infuse-factor>
2. ADYNOVATE. Prescribing Information, US FDA. Revised 06/2021. Accessed from: <https://www.fda.gov/media/94470/download>

DISCLAIMERS: The information contained herein is available to the public for information purposes only; it should not be used for diagnosing or treating a health problem or disease. It is not intended to substitute for consultation with a healthcare provider. Please consult your healthcare provider for further advice, diagnosis or treatment. The information provided is aimed at increasing awareness on Hemophilia and it is not meant for promotion of any specific product. This material contains copyright protected information, content; the use of which is limited by law and this material cannot be reproduced, replicated, used or modified by any person for their own use or further distribution. Takeda makes no representation or warranty concerning the information / content provided.

Takeda and Takeda logo are trademark or registered trademark of Takeda Pharmaceutical Company Limited.

Purna Samparq is a trademark of Takeda Biopharmaceuticals India Pvt. Ltd (formerly known as Baxalta Bioscience India Pvt. Ltd).

“Copyright © 2023 Takeda Pharmaceutical Company Limited. All rights reserved.” All trademarks are the property of their respective owners.

CONTACT US: Takeda Biopharmaceuticals India Pvt. Ltd.
(formerly known as Baxalta Bioscience India Pvt. Ltd.)

6th Floor, Tower C, Building No. 8, DLF Cyber City, DLF Phase II, Gurgaon, Haryana- 122001, India.
Tel: +91-124-455 9100 | www.takeda.com